

## SoftShield Nitrile Powder Free Examination Gloves



Order No.	Size
241-45A	SMALL
242-45A	MEDIUM
243-45A	LARGE
244-45A	X-LARGE

- 100% Nitrile contains no plasticizers, silicone, rubber, powder, or protein, reducing the potential for allergic reaction due to latex allergies.
- Nitrile Gloves can be used to promote latex free and latex safe environments.
- Effective barrier from blood borne pathogens.
- Fully textured for increased grip sensitivity-wet or dry.
- Durable with tactile sensitivity and soft feel.
- High tensile strength creates less chance of tearing.
- Beaded cuff allows for easy donning of glove.
- Extra soft, stretchy glove reduces hand fatigue.
- 9 1/2" length, ambidextrous gloves fit either hand.
- Cobalt Blue.
- Layer packed for easy dispensing, reducing wrinkles and clumping.
- 100 gloves per box, 10 boxes per case, 1,000 gloves per case, sizes S, M, L, X-L.
- Meets ASTM D6319
- Complies with FFDCA for food contact.
- Meets ASTM D6978-05 for Chemotherapy Drugs listed on the attached chart.

## **SoftShield Nitrile Powder Free Examination Gloves**

### **Chemotherapy Drug Permeation Results**

<b>CHEMOTHERAPY DRUG</b>	<b>MINIMUM BREAKTHROUGH DETECTION TIME (Minutes)</b>
Carmustine (BCNU), 3.3 mg/ml *	22.8
Thiotepa, 10.0 mg/ml *	54.6
Cisplatin, 1.0 mg/ml	No breakthrough up to 240 minutes
Cyclophosphamide (Cytoxan), 20.0 mg/ml	No breakthrough up to 240 minutes
Dacarbazine (DTIC), 10.0 mg/ml	No breakthrough up to 240 minutes
Doxorubicin Hydrochloride, 2.0 mg/ml	No breakthrough up to 240 minutes
Etoposide (Toposar), 20.0 mg/ml	No breakthrough up to 240 minutes
Fluorouracil, 50.0 mg/ml	No breakthrough up to 240 minutes
Paclitaxel (Taxol), 6.0 mg/ml	No breakthrough up to 240 minutes

Caution: Gloves used for protection against Chemotherapy drug exposure must be selected specifically for the type of chemicals used. Review material safety data sheets for the drugs being used to determine the required level of protection. \*Warning: Do not use with Carmustine (BCNU) and Thiotepa.